

Post Chemotherapy $\leq 9$ years of age	Post Chemotherapy $\geq 10$ years of age	Post HSCT $\leq 9$ years of age	Post HSCT $\geq 10$ years of age	Glossary
<p><b>Booster doses</b> for children who have received their primary course of vaccination before diagnosis.</p> <p>Please Note: These guidelines apply to children who are &lt; 2 yrs post treatment</p> <p><b>Booster dose/s commence 6 months after chemotherapy and if underlying illness is in remission.</b></p> <ul style="list-style-type: none"> <li>• <b>Infanrix Hexa</b></li> <li>• <b>Prevenar 13</b></li> <li>• <b>Nimenrix/Menveo</b></li> <li>• <b>Pneumovax 23</b> <ul style="list-style-type: none"> <li>○ minimum 8 weeks post <i>Prevenar 13</i></li> <li>○ consider if <math>\geq 2</math> years of age</li> </ul> </li> </ul>	<p><b>Booster doses</b> for children who have received their primary course of vaccination before diagnosis.</p> <p>Please Note: These guidelines apply to children who are &lt; 2 yrs post treatment</p> <p><b>Booster dose/s commence 6 months after chemotherapy and if underlying illness is in remission.</b></p> <ul style="list-style-type: none"> <li>• <b>Boostrix</b></li> <li>• <b>IPOL</b></li> <li>• <b>Prevenar 13</b></li> <li>• <b>Nimenrix/Menveo</b></li> <li>• <b>Hepatitis B</b> <ul style="list-style-type: none"> <li>○ Paediatric dose <math>\leq 10</math> years</li> <li>○ Adult dose <math>\geq 11</math> years</li> </ul> </li> <li>• <b>Gardasil 9</b> <ul style="list-style-type: none"> <li>○ Age <math>\geq 12</math> years - <math>\leq 14</math> years give 2 doses (0, and 6 – 12 months)</li> <li>○ Age <math>\geq 15</math> years give 3 doses (0, 2 and 6 months)</li> </ul> </li> <li>• <b>Pneumovax 23</b> <ul style="list-style-type: none"> <li>○ minimum 8 weeks post <i>Prevenar 13</i></li> </ul> </li> </ul>	<p>Please Note: These guidelines apply to children who are &lt; 2 yrs post treatment</p> <p><b>6 Months post HSCT (Visit 1)</b></p> <ul style="list-style-type: none"> <li>• <b>Prevenar 13 – dose 1</b></li> <li>• <b>Infanrix Hexa – dose 1</b></li> <li>• <b>Nimenrix/Menveo – dose 1</b></li> <li>• <b>Bexsero - dose 1</b></li> </ul> <p><b>Visit 2 (8 weeks post visit 1)</b></p> <ul style="list-style-type: none"> <li>• <b>Prevenar 13 – dose 2</b></li> <li>• <b>Infanrix Hexa – dose 2</b></li> <li>• <b>Nimenrix/Menveo – dose 2</b></li> <li>• <b>Bexsero - dose 2</b></li> </ul> <p><b>Visit 3 (8 weeks post visit 2)</b></p> <ul style="list-style-type: none"> <li>• <b>Prevenar 13 – dose 3</b></li> <li>• <b>Infanrix Hexa – dose 3</b></li> </ul> <p><b>Visit 4 (8 weeks post visit 3)</b></p> <ul style="list-style-type: none"> <li>• <b>Pneumovax 23</b></li> </ul>	<p>Please Note: These guidelines apply to children who are &lt; 2 yrs post treatment</p> <p><b>6 Months post HSCT (Visit 1)</b></p> <ul style="list-style-type: none"> <li>• <b>Prevenar 13 – dose 1</b></li> <li>• <b>Boostrix – dose 1</b></li> <li>• <b>IPOL – dose 1</b></li> <li>• <b>ActHIB – dose 1</b></li> <li>• <b>Hepatitis B (Adult) – dose 1</b></li> <li>• <b>Nimenrix/Menveo – dose 1</b></li> <li>• <b>Bexsero – dose 1</b></li> <li>• <b>Gardasil 9 - dose 1 ( <math>\geq 12</math> yrs of age )</b></li> </ul> <p><b>Visit 2 (8 weeks post visit 1)</b></p> <ul style="list-style-type: none"> <li>• <b>Prevenar 13 – dose 2</b></li> <li>• <b>Boostrix – dose 2</b></li> <li>• <b>IPOL – dose 2</b></li> <li>• <b>ActHIB – dose 2</b></li> <li>• <b>Hepatitis B (Adult) – dose 2</b></li> <li>• <b>Nimenrix/Menveo – dose 2</b></li> <li>• <b>Bexsero – dose 2</b></li> <li>• <b>Gardasil 9- dose 2 ( <math>\geq 12</math> yrs of age )</b></li> </ul> <p><b>Visit 3 (8 weeks post visit2)</b></p> <ul style="list-style-type: none"> <li>• <b>Prevenar 13 – dose 3</b></li> <li>• <b>Boostrix – dose 3</b></li> <li>• <b>IPOL – dose 3</b></li> <li>• <b>ActHIB – dose 3</b></li> <li>• <b>Hepatitis B (Adult) – dose 3</b></li> </ul> <p><b>Visit 4 (8 weeks post visit 3)</b></p> <ul style="list-style-type: none"> <li>• <b>Pneumovax 23</b></li> <li>• <b>Gardasil 9 – dose 3 ( <math>\geq 12</math> yrs of age )</b></li> </ul>	<p><u>ActHIB</u> * Haemophilus influenza type B</p> <p><u>Bexsero</u> * Meningococcal B</p> <p><u>Boostrix</u> * Diphtheria * Tetanus * Pertussis</p> <p><u>IPOL</u> * Inactivated Polio</p> <p><u>Gardasil 9</u> * Human Papillomavirus</p> <p><u>Hepatitis B</u> * Adult dose or paediatric dose</p> <p><u>Infanrix Hexa</u> * Diphtheria * Tetanus * Pertussis * Polio * Hib * Hepatitis B (paed)</p> <p><u>Nimenrix/Menveo</u> * Meningococcal ACWY</p> <p><u>Pneumovax 23</u> * Pneumococcal Polysaccharide</p> <p><u>Prevenar 13</u> * Pneumococcal Conjugate</p>
<p><b>Post Chemotherapy Live vaccines</b></p> <p><b>Only give on consultation with oncologist</b></p> <ul style="list-style-type: none"> <li>• <b>Measles, Mumps and Rubella</b> 2 Doses</li> <li>• <b>Varicella</b> 2 Doses</li> </ul> <p><u>Please note</u></p> <ul style="list-style-type: none"> <li>• Live vaccines must be given on the same day or 4 weeks apart</li> <li>• Do not use combination Measles, Mumps and Rubella (MMRV) vaccine in children &lt;4 yrs of age for the 1<sup>st</sup> measles vaccine</li> <li>• If child was given MMR prior to tx, MMRV <u>can</u> be used as dose 1 booster</li> <li>• MMRV may be used as the 2<sup>nd</sup> dose for all ages</li> </ul>	<p><b>Post Chemotherapy Live vaccines</b></p> <p><b>Only give on consultation with oncologist</b></p> <ul style="list-style-type: none"> <li>• <b>Measles, Mumps and Rubella</b> 2 doses</li> <li>• <b>Varicella</b> 2 doses</li> </ul> <p><u>Please note</u></p> <ul style="list-style-type: none"> <li>• Combination MMRV vaccine may be used in this group for both doses</li> </ul>	<p><b>Post HSCT Live vaccines</b></p> <p><b>Only give on consultation with oncologist</b></p> <ul style="list-style-type: none"> <li>• <b>Measles, Mumps and Rubella</b> 2 doses</li> <li>• <b>Varicella</b> 2 doses</li> </ul> <p><u>Please note</u></p> <ul style="list-style-type: none"> <li>• Live vaccines must be given on the same day or 4 weeks apart</li> <li>• Do not use combination Measles, Mumps and Rubella (MMRV) vaccine in children &lt;4 yrs of age for the 1<sup>st</sup> measles vaccine</li> <li>• MMRV may be used as the 2<sup>nd</sup> dose for all ages</li> </ul>	<p><b>Post HSCT Live vaccines</b></p> <p><b>Only give on consultation with oncologist</b></p> <ul style="list-style-type: none"> <li>• <b>Measles, Mumps and Rubella</b> 2 doses</li> <li>• <b>Varicella</b> 2 doses</li> </ul> <p><u>Please note</u></p> <ul style="list-style-type: none"> <li>• Combination MMRV vaccine may be used in this group for both doses</li> </ul>	<p><u>Measles, Mumps and Rubella</u> * Priorix * MMRII</p> <p><u>Measles, Mumps, Rubella, Varicella</u> * Priorix -tetra * ProQuad</p> <p><u>Varicella</u> * Varivax * Varilrix</p>
<p><b>Influenza</b>- Recommend annually if <math>\geq 6</math>months. 2 doses (4 weeks apart) required in the first year of receiving the vaccine <a href="http://www.mvec.vic.edu.au/immunisation-references/influenza-vaccine-recommendations/">http://www.mvec.vic.edu.au/immunisation-references/influenza-vaccine-recommendations/</a> If <b>splenectomy/hyposplenism</b> refer to MVEC for asplenia guideline <a href="http://www.mvec.vic.edu.au/immunisation-references/asplenia/">http://www.mvec.vic.edu.au/immunisation-references/asplenia/</a></p>				